

GL-TR-89-0087

AD-A208 711

Differential Geodesy of the Eotvos  
Torsion Balance

J. D. Zund

New Mexico State University  
Department of Mathematical Sciences  
Las Cruces, NM 88003-0001

April 26, 1989

Final Report  
June 24, 1986-June 23, 1988

SDTICD  
ELECTE  
JUN 09 1989  
H


APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED

GEOPHYSICS LABORATORY  
AIR FORCE SYSTEMS COMMAND  
UNITED STATES AIR FORCE  
TRANSOM AIR FORCE BASE, MASSACHUSETTS 01731-5000


89 6 09 067

This technical report has been reviewed and is approved for publication.

  
CHRISTOPHER JEKELI  
Contract Manager

  
THOMAS P. ROONEY, Chief  
Geodesy & Gravity Branch

FOR THE COMMANDER

  
DONALD H. ECKHART, Director  
Earth Sciences Division

This report has been reviewed by the ESD Public Affairs Office (PA) and is releasable to the National Technical Information Service (NTIS).

Qualified requestors may obtain additional copies from the Defense Technical Information Center. All others should apply to the National Technical Information Service.

If your address as changed, or if you wish to be removed from the mailing list, or if the addressee is no longer employed by your organization, please notify AFGL/DAA, Hanscom AFB, MA 01731-5000. This will assist us in maintaining a current mailing list.

Do not return copies of this report unless contractual obligations or notices on a specific document requires that it be returned.

## REPORT DOCUMENTATION PAGE

1a REPORT SECURITY CLASSIFICATION Unclassified			1b RESTRICTIVE MARKINGS	
2a SECURITY CLASSIFICATION AUTHORITY			3 DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; Distribution unlimited	
2b DECLASSIFICATION/DOWNGRADING SCHEDULE				
4 PERFORMING ORGANIZATION REPORT NUMBER(S)			5 MONITORING ORGANIZATION REPORT NUMBER(S) GL-TR-89-0087	
5a NAME OF PERFORMING ORGANIZATION New Mexico State University		6b OFFICE SYMBOL (if applicable)	7a NAME OF MONITORING ORGANIZATION Geophysics Laboratory	
6c ADDRESS (City, State, and ZIP Code) Department of Mathematical Sciences Las Cruces, NM 88003-0001			7b ADDRESS (City, State, and ZIP Code) Hanscom AFB Massachusetts 01731-5000	
8a NAME OF FUNDING SPONSORING ORGANIZATION		8b OFFICE SYMBOL (if applicable)	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F19628-86-K-0028	
8c ADDRESS (City, State, and ZIP Code)			10 SOURCE OF FUNDING NUMBERS	
			PROGRAM ELEMENT NO 61102F	PROJECT NO 2309
			TASK NO G1	WORK UNIT ACCESSION NO. BQ
11 TITLE (Include Security Classification) Differential Geodesy of the Eotvos Torsion Balance				
12 PERSONAL AUTHOR(S) J. D. Zund				
13a TYPE OF REPORT FINAL REPORT		13b TIME COVERED FROM 6/24/86 TO 6/23/88	14 DATE OF REPORT (Year, Month, Day) 1989 April 26	15 PAGE COUNT 16
16 SUPPLEMENTARY NOTATION				
17 COSATI CODES			18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	Differential geodesy	
			Hotine formalism	
			Curvatures	
			Eotvos torsion balance	
19 ABSTRACT (Continue on reverse if necessary and identify by block number)				
<p>The differential geodesy of the Eotvos torsion balance is examined by using the fourth and fifth fundamental tensors of the geopotential surfaces of a rotating Earth. It is shown that both these tensors involve curvature differences which are measurable by the Eotvos torsion balance. This suggests that the inability to determine the individual curvatures is not only a physical limitation, but also a geometric one. The fundamental tensors are expressed using the Hotine 2-leg formalism and their components are explicitly exhibited for the case of a spheroid. /</p>				
20 DISTRIBUTION STATEMENT OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21 ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a NAME OF RESPONSIBLE INDIVIDUAL Christopher Jekeli			22b TELEPHONE (Include Area Code)	22c OFFICE SYMBOL GL/LWG

## CONTENTS

	<u>Page</u>
1. Introduction	1
2. The Fourth & Fifth Fundamental Tensors and Invariant Curvatures	3
3. Hotine's 2-Leg Representation	5
4. Appendix	10
Acknowledgements	11
References	12



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

## 1. Introduction

Although torsion balance measurements are obsolete in gravity prospecting, the Eötvös torsion balance remains a useful conceptual device in theoretical geodesy. Indeed it may be employed to measure the curvature differences and gravity gradients which are essentially the components of the Marussi tensor of geopotential gravity gradients.

Following Hotine (1969) let  $N$  denote the geopotential function for the Earth which is assumed to be rotating with a constant angular velocity  $\tilde{\omega}$ . Points having a common value of  $N$ , say a constant  $C$ , are said to lie on a geopotential  $N$ -surface,  $N(x^1, x^2, x^3) = C$ , where  $x^r$  are rectilinear Cartesian coordinates in Euclidean 3-space. The gradient of  $N$ , denoted by  $N_r$ , satisfies the basic gradient equation

$$N_r = n v_r \quad (1.1)$$

where  $n$  is the magnitude of the acceleration due to gravity, and  $v_r$  is an upward pointing unit normal to the  $N$ -surface. Thus, although  $N_r$  is the gravity vector of the Earth, by Hotine's convention in order to give it a downward pointing direction it is necessary to take  $n = -g$ . We will simply retain  $n$  in our discussion. The gravity gradients are then given by the components of the Marussi tensor, i.e.

$$N_{rs} = N_{r,s} = n_{,s} v_r + n v_{r,s} \quad (1.2)$$

where the comma denotes covariant differentiation with respect to the flat space metric of Euclidean 3-space. The external geopotential field  $N$  of the Earth then satisfies the field equation

$$\Delta N = -2\tilde{\omega}^2, \quad (1.3)$$

where  $\Delta$  is the flat 3-dimensional Laplacian operator, and this together with the Bruns equation

$$(\log n)_{,r} v^r = -v^r_{,r} - 2\tilde{\omega}^2/n \quad (1.4)$$

reduces the number of independent components of the (symmetric) Marussi tensor from six to five. Thus, as Marussi (1949) showed, the metrical properties of the Earth's geopotential field are completely described by five functions which Hotine (1969) denoted by  $k_1, k_2, t_1, \gamma_1, \gamma_2$  and called the curvature parameters. An explicit identification of these curvature parameters will be given in Section 3.

An immediate question arises as to how in practice these curvature parameters can be measured. Both Marussi and Hotine suggested that this could be done by employing the Eötvös torsion balance and the latter worked out the corresponding theory in Hotine (1957, pages 9-10) and Hotine (1969, pages 150-151). In the latter reference he commented that

"Measurement in several azimuths will accordingly determine  $k_1 - k_2, t_1, \gamma_1, \gamma_2$  and some instrument constants, but will not separate  $k_1$  and  $k_2$ . To do this, we need an additional form of measurement."

Hotine then discussed possibilities for more refined measurements and gave several references. The general theory was analyzed exhaustively in Mueller (1960), and Mueller (1963). The general conclusion is that the inability to distinguish, i.e. measure,  $k_1$  and  $k_2$  separately is a physical limitation imposed by the use of the Eötvös torsion balance.

In the present paper we will show by employing some tensorial methods recently developed by the author in Zund (1988a, 1988b) that the average curvature differences  $\pm(k_2 - k_1)/2$  occur in expressions for the fourth and fifth fundamental tensors. This suggests that the above limitation may have a geometric origin since these tensors naturally arise in the differential geometry of the N-surfaces. In Section 2 we briefly review the relevant material from our papers which is required in this study. These results are reformulated in the 2-leg representation in Section 3, and finally in an

Appendix we exhibit an explicit 2-leg representation when the N-surfaces are spheroids.

In our discussion, unless specified to the contrary, our notation and terminology follows that employed in Hotine (1969), although this is slightly different from the purely mathematical presentation in Zund (1988a, 1988b). Special note should also be made of the elegant expose of the differential geometry of the gravity field recently given by E.W. Grafarend (1986) in this journal.

## 2. The Fourth and Fifth Fundamental Tensors and Invariant Curvatures.

It is well known that the intrinsic and extrinsic differential geometry of surfaces which are isometrically imbedded in Euclidean 3-space can be described by *three fundamental tensors*:  $a_{\alpha\beta}$ ,  $b_{\alpha\beta}$ ,  $c_{\alpha\beta}$ ; and two invariant curvatures: the Gauss (*total*) curvature  $K$  and the Germain (*mean*) curvature  $H$ . However, one can also introduce a fourth and fifth fundamental tensor and two additional invariant curvatures. These quantities are implicit in the work of the Soviet school of tensor analysis created by V.F. Kagan (1869-1953). Unfortunately, much of this work is little known or available in the West, but recently in Zund (1988a, 1988b) we have presented a unified exposition and generalization of some of this work. We now review those results which are useful in the differential geodesy of the Eötvös torsion balance.

We begin with the Euler tensor

$$h_{\alpha\beta} = \epsilon^{\rho\sigma} a_{\alpha\rho} b_{\beta\sigma} \quad (2.1)$$

which occurs in the equation

$$h_{\alpha\beta} \lambda^{\alpha} \lambda^{\beta} = 0 \quad (2.2)$$

which defines the principal directions  $\lambda^a$  on a surface, see McConnell (1931, page 212). The Euler tensor (2.1) has no privileged symmetry, although only its symmetric part contributes to defining the principal directions. If this tensor is decomposed into its symmetric and skew-symmetric parts, then we obtain

$$h_{\alpha\beta} = d_{\alpha\beta} + H\epsilon_{\alpha\beta}$$

where  $d_{\alpha\beta} = d_{\beta\alpha}$  is the fourth fundamental tensor. It was employed by S.G. Gasparyan (1961) in a beautiful memoir on the bending of surfaces which leaves the Gaussian curvature invariant, however, we believe it was known many years earlier by geometers of the Moscow school. Hence, it is probably appropriate to call it the Kagan tensor. One can also formally introduce a fifth fundamental tensor  $f_{\alpha\beta}$  which can be expressed in the form

$$f_{\alpha\beta} = H a_{\alpha\beta} - b_{\alpha\beta} . \quad (2.4)$$

Note that  $f_{\alpha\beta}$  is obviously symmetric, and by virtue of its interpretation in differential geodesy given in Section 3 we call it the Eötvös tensor. The detailed geometric and algebraic properties of these tensors are given in Zund (1988a, 1988b).

The two additional invariant curvatures are defined by

$$E = H^2 - K , \quad (2.5)$$

$$F = 2H^2 - K , \quad (2.6)$$

and are called the Euler and Monge curvatures respectively. Up to a numerical factor of one half,  $F$  is the Casorati curvature of Marussi (1951; 1985 page 19). Each of the invariant curvatures can be expressed tensorially, i.e.

$$2H = \epsilon^{\alpha\rho} \epsilon^{\beta\sigma} a_{\alpha\beta} b_{\rho\sigma} , \quad (2.7)$$



$$2K = \epsilon^{\alpha\rho} \epsilon^{\beta\sigma} b_{\alpha\beta} b_{\rho\sigma} , \quad (2.8)$$

$$2E = a^{\alpha\rho} a^{\beta\sigma} d_{\alpha\beta} d_{\rho\sigma} , \quad (2.9)$$

$$2F = a^{\alpha\rho} a^{\beta\sigma} b_{\alpha\beta} b_{\rho\sigma} . \quad (2.10)$$

An equivalent expression for (2.7) is  $2H = a^{\alpha\beta} b_{\alpha\beta}$  which shows that the Eötvös tensor is traceless.

### 3. Hotine's 2-Leg Representation.

Following Grafarend (1986) we call a pair of orthogonal unit surface vectors a 2-leg. In accord with the notation of Chapters 6 and 7 of Hotine (1969), we denote the contravariant components of the 2-leg by  $\{\rho^\alpha, j^\alpha\}$  with  $\rho^r := x_\alpha^r \rho^\alpha$ ,  $j^r := x_\gamma^r j^\gamma$  being the corresponding space vectors. In the latter expressions,  $x_\alpha^r := \partial x^r / \partial x^\alpha$  where the  $x^r$  are rectangular Cartesian coordinates in Euclidean 3-space, and  $x^\alpha$  denote curvilinear coordinates, i.e. parameters, on the N-surfaces. This 2-leg may be extended to a 3-leg of space vectors  $\{\rho^r, j^r, v^r\}$  by including the unit normal  $v^r$  to the N-surfaces. Then the 2-legs  $\{\rho^r, v^r\}$ ,  $\{j^r, v^r\}$  define planes which are said to be sections of the N-surfaces.

As a consequence of the orthogonality conditions of the 2-leg, the Levi-Civita permutation tensor has the 2-leg representation

$$\epsilon_{\alpha\beta} = \rho_\alpha j_\beta - j_\alpha \rho_\beta . \quad (3.1)$$

The fundamental surface tensors are then readily found to have the representations:

$$a_{\alpha\beta} = \rho_\alpha \rho_\beta + j_\alpha j_\beta , \quad (3.2)$$

$$b_{\alpha\beta} = k_1 \rho_\alpha \rho_\beta + l_1 (\rho_\alpha j_\beta + j_\alpha \rho_\beta) + k_2 j_\alpha j_\beta , \quad (3.3)$$

$$c_{\alpha\beta} = k_1^2 \rho_\alpha \rho_\beta + (k_1 + k_2) t_1 (\rho_\alpha j_\beta + j_\alpha \rho_\beta) + k_2^2 j_\alpha j_\beta, \quad (3.4)$$

where  $k_1, k_2$  are the normal curvatures, and  $t_1, t_2 = -t_1$  are the geodesic torsions of the N-surface in the directions  $\rho^\alpha, j^\alpha$ . The remaining two curvature parameters  $\gamma_1, \gamma_2$  are related to the curvature of the normal  $\rho^\alpha$  in the directions  $\rho^\alpha, j^\alpha$  and do not depend on the fundamental tensors.

The corresponding expressions for the invariant curvatures of the N-surfaces are given by

$$H = (k_1 + k_2)/2, \quad (3.5)$$

$$K = k_1 k_2 - t_1^2, \quad (3.6)$$

$$E = (k_1 - k_2)^2/4 + t_1^2, \quad (3.7)$$

$$F = (k_1^2 + k_2^2)/2 + t_1^2. \quad (3.8)$$

The 2-leg representations for the Kagan and Eötvös tensors are then found to be

$$d_{\alpha\beta} = t_1 \rho_\alpha \rho_\beta + \left(\frac{k_2 - k_1}{2}\right) (\rho_\alpha j_\beta + j_\alpha \rho_\beta) + t_1 j_\alpha j_\beta, \quad (3.9)$$

$$f_{\alpha\beta} = \left(\frac{k_2 - k_1}{2}\right) (\rho_\alpha \rho_\beta - j_\alpha j_\beta) - t_1 (\rho_\alpha j_\beta + j_\alpha \rho_\beta), \quad (3.10)$$

and hence the Euler tensor has the representation

$$h_{\alpha\beta} = t_1 (\rho_\alpha \rho_\beta + j_\alpha j_\beta) + k_2 \rho_\alpha j_\beta - k_1 j_\alpha \rho_\beta. \quad (3.11)$$

Note that these representations show that in general neither  $d_{\alpha\beta}$  nor  $h_{\alpha\beta}$  are traceless, viz both have the trace  $2t_1$ , while as mentioned previously  $f_{\alpha\beta}$  is always traceless.

Equations (3.9), (3.10) indicate the first evidence of the relationship between the curvature differences and the Kagan and Eötvös tensors. This

appearance is of course due to our choice of the 2-leg, but aside from the orthogonality requirement the 2-leg  $\{\rho_\alpha, j_\alpha\}$  is quite general. Indeed, if this requirement were relaxed, and the unit vectors  $\rho^\alpha, j^\alpha$  were inclined to each other at an angle  $0 < \theta < \pi$ , then  $a_{\alpha\beta} \rho^\alpha j^\beta = \cos \theta$ ,  $\epsilon_{\alpha\beta} \rho^\alpha j^\beta = \sin \theta$ , and our previous expressions would contain additional trigonometric functions but no new curvature parameters would occur. Thus the relationship between the curvature differences and these tensors is not essentially a consequence of a specialization of the 2-leg.

However, by a further natural specialization of the 2-leg, this representation can be put in a more transparent and striking form. This additional specialization involves choosing a 2-leg in which the normal curvatures  $k_1, k_2$  assume their maximum and minimum values. Such a 2-leg is called a *principal* 2-leg, the corresponding vectors are *principal directions*, and the sections of the N-surfaces *principal sections*. Following Hotine (1969, page 42) these vectors are denoted by  $u^\alpha, v^\alpha$  and relative to this 2-leg,  $k_1, k_2$  reduce to the *principal curvatures*  $\kappa_1, \kappa_2$ , and  $t_1 = 0$ . Algebraically this specialization makes  $u^\alpha, v^\alpha$  *eigenvectors* of the fundamental tensors  $a_{\alpha\beta}, b_{\alpha\beta}, c_{\alpha\beta}, d_{\alpha\beta}$ , and  $f_{\alpha\beta}$ . The principal 2-leg representations of these tensors are thus:

$$a_{\alpha\beta} = u_\alpha u_\beta + v_\alpha v_\beta, \quad (3.12)$$

$$b_{\alpha\beta} = \kappa_1 u_\alpha u_\beta + \kappa_2 v_\alpha v_\beta, \quad (3.13)$$

$$c_{\alpha\beta} = \kappa_1^2 u_\alpha u_\beta + \kappa_2^2 v_\alpha v_\beta, \quad (3.14)$$

$$d_{\alpha\beta} = \left(\frac{\kappa_2 - \kappa_1}{2}\right) (u_\alpha v_\beta + v_\alpha u_\beta), \quad (3.15)$$

$$f_{\alpha\beta} = \left(\frac{\kappa_2 - \kappa_1}{2}\right) u_\alpha u_\beta + \left(\frac{\kappa_1 - \kappa_2}{2}\right) v_\alpha v_\beta, \quad (3.16)$$

and this representation explicitly exhibits the *eigenvalues* of these tensors in terms of the principal curvatures. Of course, this *eigenvector-eigenvalue* property does not hold for either the Levi-Civita tensor

$$e_{\alpha\beta} = u_{\alpha} v_{\beta} - v_{\alpha} u_{\beta} , \quad (3.17)$$

or the Euler tensor

$$h_{\alpha\beta} = \kappa_2 u_{\alpha} v_{\beta} - \kappa_1 v_{\alpha} u_{\beta} . \quad (3.18)$$

Higher order powers of the principal curvatures may occur, for example in

$$2Hc_{\alpha\beta} - Kb_{\alpha\beta} = \kappa_1^3 u_{\alpha} u_{\beta} + \kappa_2^3 v_{\alpha} v_{\beta} ,$$

but the analysis given in Zund (1988b) suggests that such combinations of the fundamental tensors have no particular geometric significance.

Although the Kagan tensor (3.15) involves a curvature difference, the Eötvös tensor (3.16) is more interesting since in terms of its principal 2-leg representation it exhibits *both* curvature differences. Indeed, this is why we refer to this tensor as the Eötvös tensor. It should be noted that with characteristic modesty Marussi (1931, 1985) always referred to  $N_{rs}$ , i.e. our (1.2), as the Eötvös tensor, but most mathematical geodesists have followed Hotine's custom of calling the latter the Marussi tensor. Note that since  $\kappa_1 = \kappa_2$  for a sphere, the Kagan and Eötvös tensors in differential geodesy essentially measure the deviation of the spheroidal N-surfaces from perfect spherical form. The appearance of the curvature differences in these tensors is surprising since both tensors have a purely geometric origin. It also suggests that Hotine's observation quoted in Section 1 is not merely a physical limitation imposed by employing torsion balance measurements, but a geometric limitation arising from the geometry of the N-surfaces. An additional reason for this can be found in Hotine (1969, p. 44) where it is

shown that the principal 2-leg representation of the Codazzi-Mainardi equations leads to the equations

$$(\kappa_1 - \kappa_2)\sigma = \kappa_{1,\alpha} v^\alpha$$

$$(\kappa_1 - \kappa_2)\sigma^* = \kappa_{2,\alpha} u^\alpha$$

where  $\sigma, \sigma^*$  are the geodesic curvatures of the principal directions  $u^\alpha, v^\alpha$  respectively.

An excellent classical discussion, without using tensors, is given in Slotnick (1932), and it is reviewed with reference to geophysical prospecting in Dobrin (1960, pp. 180-183).

We believe that the above discussion indicates not only a connection between the torsion balance and differential geometry, but also the advantages of the 2-leg approach to differential geodesy. This approach was implicit in the work of Marussi and Hotine, however, their exposition of it failed to emphasize that it is a *coordinate-free method* and that it is conceptually distinct from the classical coordinate approach to tensors. In the 2-leg approach, as we will illustrate in the following Appendix, the rôle of coordinates is a secondary one and the resulting coordinates have a natural interpretation. In effect one makes a physical, or geometrical, choice of a 2-leg, which involves measurable quantities and then introduces a coordinate system based on the 2-leg. This is of course classically the notion of *physical components* of vectors/tensors as contained in the appendix of McConnell (1931), however, only the modern approach to tensors makes this procedure natural. Both Marussi and Hotine realized the advantages of the 2-leg formalism, but were unable to separate it from their primary goal of finding *intrinsic* coordinate systems.

#### 4. Appendix. Explicit Calculations on the Spheroid.

We now exhibit the components of the fundamental tensor, the Euler tensor, and the principal 2-legs when the N-surfaces are spheroids, i.e., ellipsoids of revolution about the  $x^3$ -axis having semi-axes  $a > b$ . Following Hotine (1969), we take the curvilinear surface coordinates (parameters) to be  $x^\alpha = (\omega, \vartheta)$  where  $\omega$  is the longitude and  $\vartheta$  the latitude  $0 \leq \omega < 2\pi$ ,  $-\pi/2 \leq \vartheta \leq \pi/2$ . Then the first fundamental form becomes

$$ds^2 = \rho_1^2 \cos^2 \vartheta d\omega^2 + \rho_2^2 d\vartheta^2 \quad (\text{A.1})$$

where  $\rho_1 = 1/\kappa_1$ ,  $\rho_2 = 1/\kappa_2$ , are the principal radii of curvature where

$$\kappa_1 = -(1 - e^2 \sin^2 \vartheta)^{1/2}/a, \quad (\text{A.2})$$

$$\kappa_2 = -(1 - e^2 \sin^2 \vartheta)^{3/2}/a(1 - e^2), \quad (\text{A.3})$$

and  $e$  is the eccentricity

$$e^2 = (a^2 - b^2)/a^2. \quad (\text{A.4})$$

In the remainder of our discussion it will be convenient to employ  $\kappa_1$ ,  $\kappa_2$ , or  $\rho_1$ ,  $\rho_2$  in place of the above explicit values. Then the components of the first fundamental tensor are given by

$$\|a_{\alpha\beta}\| = \left\| \begin{array}{cc} \rho_1^2 \cos^2 \vartheta & , & 0 \\ 0 & , & \rho_2^2 \end{array} \right\| \quad (\text{A.5})$$

and by (3.12) the components of the principal 2-leg are

$$u_1 = -\rho_1 \cos \vartheta, \quad u_2 = 0; \quad (\text{A.6})$$

$$v_1 = 0, \quad v_2 = -\rho_2; \quad (\text{A.7})$$

and

$$u^1 = -\kappa_1 \sec \vartheta, \quad u^2 = 0; \quad (\text{A.8})$$

$$v^1 = 0, v^2 = -\kappa_2. \quad (\text{A.9})$$

The components of the remaining fundamental tensors are then

$$\|b_{\alpha\beta}\| = \begin{vmatrix} \kappa_1 \rho_1^2 \cos^2 \vartheta & 0 \\ 0 & \kappa_2 \rho_2^2 \end{vmatrix}; \quad (\text{A.10})$$

$$\|c_{\alpha\beta}\| = \begin{vmatrix} \kappa_1^2 \rho_1^2 \cos^2 \vartheta & 0 \\ 0 & \kappa_2^2 \rho_2^2 \end{vmatrix}; \quad (\text{A.11})$$

$$\|d_{\alpha\beta}\| = \begin{vmatrix} 0 & (\frac{\kappa_2 - \kappa_1}{2}) \rho_1 \rho_2 \cos \vartheta \\ (\frac{\kappa_2 - \kappa_1}{2}) \rho_1 \rho_2 \cos \vartheta & 0 \end{vmatrix}; \quad (\text{A.12})$$

$$\|f_{\alpha\beta}\| = \begin{vmatrix} (\frac{\kappa_2 - \kappa_1}{2}) \rho_1^2 \cos^2 \vartheta & 0 \\ 0 & (\frac{\kappa_1 - \kappa_2}{2}) \rho_2^2 \cos^2 \vartheta \end{vmatrix}; \quad (\text{A.13})$$

and the Euler tensor has components

$$\|h_{\alpha\beta}\| = \begin{vmatrix} 0 & \rho_1 \cos \vartheta \\ -\rho_2 \cos \vartheta & 0 \end{vmatrix}. \quad (\text{A.14})$$

#### Acknowledgements

This work was supported by the Air Force Geophysics Laboratory. The author is grateful to B.H. Chovitz, C. Jekeli, and J.M. Wilkes for their comments and discussion.

## REFERENCES

- Dobrin, M.B. (1960). Introduction to geophysical prospecting, McGraw Hill, New York.
- Gasparyan, S.G. (1961). Definition of the fourth fundamental tensor of a surface given its metric and mean curvature (in Russian). *Uspekhi Matematicheskikh Nauk* 16, pp. 101-108.
- Grafarend, E.W. (1986). Differential geometry of the gravity field, *Manuscripta Geodaetica* 11, pp. 29-37.
- Hotine, M. (1957). Metrical properties of the Earth's gravitational field, A.I.G. Toronto Assembly Study Group No. 1, 29 pp.
- Hotine, M. (1969). Mathematical geodesy, U.S. Department of Commerce, Washington, D.C.
- McConnell, A.J. (1931). Applications of the absolute differential calculus, Blackie & Sons, London; reprinted in 1957 as Applications of tensor analysis, Dover Publications, New York.
- Marussi, A. (1951). Fondamenti di geodesia intrinseca, *Pubb. Comm. Geodet. Ital.*, III Ser. Mem. 7, 47 pp., translated as Foundations of intrinsic geodesy in Marussi (1985) pp. 13-58.
- Marussi, A., Chiaruttini, C. (1985). The motion of a free particle and of a spherical pendulum in the microgravitational field of a gravitationally stabilized satellite in circular orbit in a central field, in Marussi (1985), pp. 179-189.
- Marussi, A. (1985). Intrinsic geodesy, translated by W.I. Reilly, Springer Verlag, Berlin.
- Mueller, I.I. (1960). The gradients of gravity and their applications in geodesy, Ph.D. dissertation, The Ohio State University, Columbus, 240 pp.
- Mueller, I.I. (1963). Geodesy and the torsion balance, *Proc. American Society of Civil Engineers, Journal of Surveying and Mapping Division*, SU 3, 89, pp. 123-155.
- Slotnick, M.M. (1932). Curvature of equipotential surfaces, *Bull. Amer. Assoc. Petroleum Geologists* 16(12), pp. 80-89; reprinted in 1947 in early geophysical papers of the Society of Exploration Geophysicists, *Soc. Exploration Geophysicists*, Tulsa, pp. 410-419.
- Zund, J.D. (1988a). Tensorial methods in classical differential geometry I: basic principles, *Tensor N.S.* 47, in press.
- Zund, J.D. (1988b). Tensorial methods in classical differential geometry II: basic surface tensors, *Tensor N.S.* 47, in press.